

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the present application:

Listing of Claims:

1. **(Currently Amended)** A method for calculating the activity of a cyclin-dependent kinase comprising the steps of:

preparing a sample from living cells;

catching the cyclin-dependent kinase in the sample by an anti-cyclin-dependent kinase antibody;

reacting adenosine 5'-O-(3-thiotriphosphate) (ATP-γS) (~~ATP-S~~) with a substrate for the cyclin-dependent kinase in presence of the cyclin-dependent kinase in order to introduce a monothiophosphate group into a serine or threonine residue of the substrate, the substrate not containing a sulfur atom;

labeling the substrate by coupling a labeling fluorophore or a labeling enzyme with a sulfur atom of the introduced monothiophosphate group;

removing the fluorophore or the enzyme not labeling the substrate from the labeled substrate;

measuring the amount of fluorescence from the labeling fluorophore labeling the substrate, or reacting the labeling enzyme labeling the substrate with a substance which generates

an optically detectable product by reaction with the labeling enzyme and optically measuring the amount of the generated product; and

calculating the activity of the cyclin-dependent kinase from the measured amount of fluorescence or the measured amount of the generated product with reference to a pre-produced reference curve.

2. **(Previously Presented)** The method according to claim 1, wherein the cyclin-dependent kinase is selected from the group consisting of CDK1, CDK2, CDK4 and CDK6.

3. **(Original)** The method according to claim 1, wherein the labeling fluorophore is a fluorescent dye.

4. **(Original)** The method according to claim 3, wherein the fluorescent dye is FITC.

5. **(Original)** A method according to claim 1, wherein the labeling enzyme is peroxidase.

6. **(Previously Presented)** A method according to any one of claims 1 to 5, wherein the cyclin-dependent kinase is CDK1 or CDK2 and the substrate is histone H1.

7. **(Previously Presented)** A method according to any one of claims 1 to 5, wherein the cyclin-dependent kinase is CDK4 or CDK6 and the substrate is Rb whose cysteine residue is substituted by alanine.

8-9. **(Cancelled)**

10. **(Currently Amended)** A method for obtaining the activity of a cyclin-dependent kinase comprising the steps of:

preparing a sample from living cells;

catching the cyclin-dependent kinase in the sample by anti-cyclin-dependent kinase antibody;

reacting adenosine 5'-O-(3-thiotriphosphate) (ATP- γ S) (~~ATP-S~~) with a substrate for the cyclin-dependent kinase in presence of the cyclin-dependent kinase in order to introduce a monothiophosphate group into a serine or threonine residue of the substrate, the substrate not containing a sulfur atom;

labeling the substrate by coupling a labeling fluorophore or a labeling enzyme with a sulfur atom of the introduced monothiophosphate group;

removing the fluorophore or the enzyme not labeling the substrate from the labeled substrate;

measuring the amount of fluorescence from the labeling fluorophore labeling the substrate, or reacting the labeling enzyme labeling the substrate with a substance which generates

Application No. 10/074,041

Docket No.: 0397-0440P

Art Unit 1642

Reply to Office Action of May 31, 2005

an optically detectable product by reaction with the labeling enzyme and optically measuring the amount of the generated product; and

obtaining the activity of the cyclin-dependent kinase from the measured amount of fluorescence or the measured amount of the generated product.